GENERAL RELATIVITY WITH A POSITIVE COSMOLOGICAL CONSTANT Λ AS A GAUGE THEORY

MARTA DUDEK
Institute of Mathematics, University of Szczecin
Wielkopolska 15, 70-451 Szczecin, Poland, EU
e-mail: marta.dudek@vp.pl

JANUSZ GARECKI
Institute of Mathematics, University of Szczecin
Wielkopolska 15, 70-451 Szczecin, Poland, EU
e-mail: garecki@wmf.univ.szczecin.pl

Received 8 March 2017

arXiv:1703.06024v1 [gr-qc] 17 Mar 2017

In the paper we show that the general relativity action (and Lagrangian) in recent Einstein-Palatini formulation is equivalent to the action (and Langrangian) of a gauge field.

We begin with a bit of information of the Einstein-Palatini (EP) action, then we present how Einstein fields equations can be derived from it. In the next section, we consider Einstein-Palatini action integral for general relativity with a positive cosmological constant Λ in terms of $\hat{F}$, the curvature of the affine connection's pulled back from de Sitter bundle to Lorentz bundle. We will see that in terms of $\hat{F}$ this action takes the form typical for a gauge field. Finally, we give a geometrical interpretation of the curvature $\hat{F}$.

Keywords: action integral, fiber bundle, connection in a principal fiber bundle and its curvature, pull-back of forms, Lie groups and their algebras.

1 Einstein-Palatini action for general relativity

In this section we would like to remind briefly Einstein- Palatini formalism for general relativity (GR).

The Einstein field equations can be derived by postulating the Einstein-Hilbert action to be the true action for GR. Albert Einstein firstly used only metric as a independent variable to do variation of this action. The connection in this approach to EH action is the metric and symmetric Levi-Civita connection. Later Einstein and Palatini proposed to take the metric and affine connection as independent variables in the action principle. This method allowed to compute the field equations for a more general metric affine connection rather than the Levi-Civita connection. Here the spacetime admitted torsion when matter Lagrangian explicite depended on connection.Thus, the Einstein-Palatini formalism gave us a powerful tool for theories of gravitation which have more general Riemann-Cartan geometry.
In this section, we are going to present Einstein-Palatini action in recent formulation and how the Einstein fields equations can be computed from it.

The Einstein-Palatini action with cosmological constant Λ in this new formulation [3] is defined as follows

$$S_{E-P} = \frac{1}{4\kappa}\int\limits_{\mathcal{D}} \Big(\vartheta^i \wedge \vartheta^j \wedge \Omega^{kl} - \frac{\Lambda}{6}\vartheta^i \wedge \vartheta^j \wedge \vartheta^k \wedge \vartheta^l\Big)\eta_{ijkl}, \tag{1}$$

where Ω is the curvature of ω (spin connection) and $\kappa = 8\pi G/c^4$. All indices take values $(0,1,2,3)$. $\mathcal{D}$ means an established 4-dimensional compact domain in spacetime. In the above formula ϑ^a denote 1-forms of the Lorentzian coreper in term of which the spacetime looks locally Minkowskian: $g = \eta_{ik}\vartheta^i \otimes \vartheta^k$, $\eta_{ik} = diag(1,-1,-1,-1)$.
η_{ijkl} is completely antysymmetric Levi-Civita pseudotensor: $\eta_{0123} = \sqrt{|g|}$, where $g := det(g_{ik})$. In a Lorentzian coreper $|g| = 1$. Spin connection ω is a general metric connection (or Levi-Civita connection) in Lorentzian coreper.

For the geometrical units $G = c = 1$ the above formula has the extended form

$$S_{E-P} = \frac{1}{32\pi}\int\limits_{\mathcal{D}} \eta_{ijkl}\vartheta^i \wedge \vartheta^j \wedge \Omega^{kl} - \frac{\Lambda}{192\pi}\int\limits_{\mathcal{D}} \eta_{ijkl}\vartheta^i \wedge \vartheta^j \wedge \vartheta^k \wedge \vartheta^l. \tag{2}$$

Let us denote by S_1 and S_2 two integrals in the above formula of the Einstein-Palatini action and compute their variations. From the integral

$$S_1 = \frac{1}{32\pi}\int\limits_{\mathcal{D}} \eta_{ijkl}\vartheta^i \wedge \vartheta^j \wedge \Omega^{kl} = \frac{1}{16\pi}\int\limits_{\mathcal{D}} \Omega^i{}_j \wedge \eta_i{}^j, \tag{3}$$

where $\eta_i{}^j$ is the 2-form defined in Appendix 1, we get

$$\delta S_1 = \frac{1}{16\pi}\int\limits_{\mathcal{D}} \delta\Big(\Omega^i{}_j \wedge \eta_i{}^j\Big). \tag{4}$$

Below we give the method of computing $\delta\Big(\Omega^i{}_j \wedge \eta_i{}^j\Big)$ in Lorentzian coreper ϑ^i with spin connection $\omega^i{}_k$.

$$\rho\delta\Big(\Omega^i{}_j \wedge \eta_i{}^j\Big) = \rho\Big(\delta\Omega^i{}_j \wedge \eta_i{}^j + \Omega^i{}_j \wedge \delta\eta_i{}^j\Big), \tag{5}$$

where ρ denotes $\frac{1}{2\kappa} = \frac{1}{16\pi}$.

Let us take the standard expressions

$$\Omega^i{}_j = d\omega^i{}_j + \omega^i{}_k \wedge \omega^k{}_j \tag{6}$$

and

$$\begin{aligned}\eta_i{}^j &= g^{jp}\eta_{ip}\\ &= \frac{1}{2}g^{jp}\vartheta^k \wedge \eta_{ipk}\\ &= \frac{1}{2}g^{jp}\vartheta^k \wedge \vartheta^l \wedge \eta_{ipkl}\\ &= \frac{1}{2}g^{jl}\eta_{ilmn}\vartheta^m \wedge \vartheta^n.\end{aligned} \tag{7}$$

η_{ip}, η_{ipk}, η_{ipkl} are forms firstly introduced by A. Trautman [11] (see Appendix 1).

Then, we have

$$\begin{aligned}&\rho\delta\Big(\Omega^i{}_j \wedge \eta_i{}^j\Big) =\\ &= \rho\Big(\delta d\omega^i{}_j + \delta\omega^i{}_k \wedge \omega^k{}_j + \omega^i{}_k \wedge \delta\omega^k{}_j\Big) \wedge \eta_i{}^j + \frac{\rho}{2}\Omega^i{}_j \wedge \delta\Big(g^{jl}\eta_{ilmn}\vartheta^m \wedge \vartheta^n\Big).\end{aligned} \tag{8}$$

Now we use the fact that δ and d comute and that $g^{il} = const$ and $\eta_{ilmn} = const$ in the orthonormal thetrad, so the subintegral expression in δS_1 has the following form

$$\rho\delta\Big(\Omega^i{}_j \wedge \eta_i{}^j\Big)$$
$$= \rho\Big[d\big(\delta\omega^i{}_j\big) \wedge \eta_i{}^j + \delta\omega^i{}_k \wedge \omega^k{}_j \wedge \eta_i{}^j + \omega^i{}_k \wedge \delta\omega^k{}_j \wedge \eta_i{}^j + \frac{1}{2}\Omega^{il} \wedge \eta_{ilmn} \wedge \delta\big(\vartheta^m \wedge \vartheta^n\big)\Big]. \quad (9)$$

Because

$$\delta\big(\vartheta^m \wedge \vartheta^n\big) = \delta\vartheta^m \wedge \vartheta^n + \vartheta^m \wedge \delta\vartheta^n \quad (10)$$

and

$$d\big(\delta\omega^i{}_j\big) \wedge \eta_i{}^j = d\big(\delta\omega^i{}_j \wedge \eta_i{}^j\big) + \delta\omega^i{}_j \wedge d\eta_i{}^j, \quad (11)$$

so

$$\rho\delta\Big(\Omega^i{}_j \wedge \eta_i{}^j\Big) =$$
$$= \rho\Big[d\big(\delta\omega^i{}_j \wedge \eta_i{}^j\big) + \delta\omega^i{}_j \wedge d\eta_i{}^j + \delta\omega^i{}_k \wedge \omega^k{}_j \wedge \eta_i{}^j + \omega^i{}_k \wedge \delta\omega^k{}_j \wedge \eta_i{}^j\Big]$$
$$+ \rho\Big[\frac{1}{2}\Omega^{il}\eta_{ilmn} \wedge \big(\delta\vartheta^m \wedge \vartheta^n + \vartheta^m \wedge \delta\vartheta^n\big)\Big]; \quad (12)$$

where $\rho = \frac{1}{2\kappa} = \frac{c^4}{16\pi G}$.

Next, we use two laws given below:

$$\omega_p \wedge \omega_q = (-1)^{pq}\omega_q \wedge \omega_p, \quad (13)$$

$$d\big(\omega_p \wedge \omega_q\big) = d\omega_p \wedge \omega_q + (-1)^p\omega_p \wedge d\omega_q \quad (14)$$

and we change some indices. Then the variation of $\rho\delta\Big(\Omega^i{}_j \wedge \eta_i{}^j\Big)$ has the form

$$\rho\delta\Big(\Omega^i{}_j \wedge \eta_i{}^j\Big) =$$
$$= \rho\Big[d\big(\delta\omega^i{}_j \wedge \eta_i{}^j\big) + \frac{1}{2}\delta\omega^i{}_j \wedge 2\big(d\eta_i{}^j + \omega^j{}_k \wedge \eta_i{}^k - \omega^k{}_i \wedge \eta_k{}^j\big) + \delta\vartheta^i \wedge \Omega^{kl}\eta_{klim} \wedge \vartheta^m\Big]$$
$$= \delta\vartheta^i \wedge \rho\Omega^{kl}\eta_{klim} \wedge \vartheta^m + \frac{1}{2}\delta\omega^i{}_j \wedge 2\rho D\eta_i{}^j + d\big(\rho\delta\omega^i{}_j \wedge \eta_i{}^j\big) \quad (15)$$

where

$$D\eta_i{}^j = d\eta_i{}^j + \omega^j{}_k \wedge \eta_i{}^k - \omega^k{}_i \wedge \eta_k{}^j. \quad (16)$$

Finally, we get the following result

$$\rho\delta\Big(\Omega^i{}_j \wedge \eta_i{}^j\Big) = \delta\vartheta^i \wedge \rho\Omega^{kl}\eta_{klim} \wedge \vartheta^m + \rho\delta\omega^i{}_j \wedge D\eta_i{}^j + d\big(\rho\delta\omega^i{}_j \wedge \eta_i{}^j\big).$$
$$= \delta\vartheta^i \wedge \rho\Omega^{kl}\eta_{kli} + \rho\delta\omega^i{}_j \wedge D\eta_i{}^j + an\ exact\ form. \quad (17)$$

The second integral from the Einstein-Palatini action

$$S_2 = \frac{-\Lambda}{192\pi}\int\limits_{\mathcal{D}} \eta_{ijkl}\vartheta^i \wedge \vartheta^j \wedge \vartheta^k \wedge \vartheta^l \quad (18)$$

has the following variation

$$\delta S_2 = \frac{-\Lambda}{192\pi}\int\limits_{\mathcal{D}} 24\delta\vartheta^i \wedge \eta_i$$
$$= \frac{-\Lambda}{8\pi}\int\limits_{\mathcal{D}} \delta\vartheta^i \wedge \eta_i. \tag{19}$$

It was computed in a similar way as the variation for the first integral S_1.

$$\delta S_2 = \frac{-\Lambda}{192\pi}\int\limits_{\mathcal{D}} \delta\Big(\eta_{ijkl}\vartheta^i \wedge \vartheta^j \wedge \vartheta^k \wedge \vartheta^l\Big)$$
$$= \frac{-\Lambda}{192\pi}\int\limits_{\mathcal{D}} \eta_{ijkl}\delta\Big(\vartheta^i \wedge \vartheta^j \wedge \vartheta^k \wedge \vartheta^l\Big) \tag{20}$$

where

$$\delta\Big(\vartheta^i \wedge \vartheta^j \wedge \vartheta^k \wedge \vartheta^l\Big) =$$
$$= \delta\vartheta^i \wedge \vartheta^j \wedge \vartheta^k \wedge \vartheta^l + v^i \wedge \delta\vartheta^j \wedge \vartheta^k \wedge \vartheta^l + \vartheta^i \wedge \vartheta^j \wedge \delta\vartheta^k \wedge \vartheta^l + \vartheta^i \wedge \vartheta^j \wedge \vartheta^k \wedge \delta\vartheta^l$$
$$= \delta\vartheta^i \wedge \vartheta^j \wedge \vartheta^k \wedge \vartheta^l - \delta\vartheta^j \wedge \vartheta^i \wedge \vartheta^k \wedge \vartheta^l + \delta\vartheta^k \wedge \vartheta^i \wedge \vartheta^j \wedge \vartheta^l - \delta\vartheta^l \wedge \vartheta^i \wedge \vartheta^j \wedge \vartheta^k \tag{21}$$

and

$$\delta\Big(\vartheta^i \wedge \vartheta^j \wedge \vartheta^k \wedge \vartheta^l\Big)\eta_{ijkl} = 4\delta\vartheta^i \wedge \eta_{ijkl}\vartheta^j \wedge \vartheta^k \wedge \vartheta^l. \tag{22}$$

Thus, the variation δS_2 finally equals

$$\delta S_2 = \frac{-\Lambda}{192\pi}\int\limits_{\mathcal{D}} 4\delta\vartheta^i \wedge \eta_{ijkl}\vartheta^j \wedge \vartheta^k \wedge \vartheta^l \tag{23}$$

The expression $\eta_{ijkl}\vartheta^j \wedge \vartheta^k \wedge \vartheta^l$ is equal $6\eta_i$ (see Appendix 1), so using this substitution we obtain as follows

$$\delta S_2 = \frac{-\Lambda}{192\pi}\int\limits_{\mathcal{D}} 24\delta\vartheta^i \wedge \eta_i$$
$$= \frac{-\Lambda}{8\pi}\int\limits_{\mathcal{D}} \delta\vartheta^i \wedge \eta_i \tag{24}$$

Joining both variations δS_1, δS_2 with the third variation δS_m of the matter action $S_m = \int\limits_{\mathcal{D}} L_{mat}(\phi^A, D\phi^A, \vartheta^i)$, which we write in the form [11]

$$\delta S_m = \delta\int\limits_{\mathcal{D}} L_{mat}(\phi^A, D\phi^A, \vartheta^i)$$
$$= \int\limits_{\mathcal{D}}\Big(\delta\vartheta^i \wedge t_i + \frac{1}{2}\delta\omega^i_j \wedge s^j_i + \delta\phi^A \wedge L_A + an\ exact\ form\Big), \tag{25}$$

we obtain for total action, gravity and matter

$$\begin{aligned}\delta S &= \delta S_1 + \delta S_2 + \delta S_m \\ &= \int\limits_{\mathcal{D}} \Big[\frac{1}{8\pi}\delta\vartheta^i \wedge \Big(\frac{1}{2}\Omega^{kl} \wedge \eta_{kli} - \Lambda\eta_i + 8\pi t_i\Big) + an\ exact\ form\Big] \\ &+ \int\limits_{\mathcal{D}} \Big[\frac{1}{2}\delta\omega^i_j \wedge \Big(\frac{1}{8\pi}D\eta^j_i + s^j_i\Big) + \delta\phi^A \wedge L_A + an\ exact\ form\Big]. \end{aligned} \tag{26}$$

The form of the variation $\delta L_m = \delta\vartheta^i \wedge t_i + \frac{1}{2}\delta\omega^i{}_j \wedge s_i{}^j + \delta\phi^A \wedge L^A + an\ exact\ form$ defines the 3-forms t_i, $s_i{}^j$, L^A of the energy-momentum classical spin and the left hand side of the equations of motion for matter respectively. In the above formulas ϕ^A means tensor-valued matter form and $D\phi^A$ its absolute exterior derivative.

Einstein's equations like all the other physical field equations arise due to variational principle, which is called the Principle of Stationary Action or Hamiltonian Principle. In our case it has the following form:

$$\delta S = \delta \int\limits_{\Omega} \Big(L_g + L_m\Big) = 0, \tag{27}$$

where variations $\delta\vartheta^i$, $\delta\omega^i{}_j$ and $\delta\phi^A$ are vanishing on the boundary $\delta\mathcal{D}$ of the compact domain $\mathcal{D}$.
Here $L_g = \frac{1}{2\kappa}\Big(\Omega^i{}_j \wedge \eta_i{}^j - \Lambda\eta\Big)$ and represents the Lagrangian density of the gravitational field and L_m is the Langrangian density of matter.

It is seen from above considerations that following Hamiltonian Principle

$$\begin{aligned}\delta S &= \int\limits_{\mathcal{D}} \Big[\frac{1}{8\pi}\delta\vartheta^i \wedge \Big(\frac{1}{2}\Omega^{kl} \wedge \eta_{kli} - \Lambda\eta_i + 8\pi t_i\Big) + an\ exact\ form\Big] \\ &+ \int\limits_{\mathcal{D}} \Big[\frac{1}{2}\delta\omega^i_j \wedge \Big(\frac{1}{8\pi}D\eta^j_i + s^j_i\Big) + \delta\phi^A \wedge L_A + an\ exact\ form\Big] \\ &= 0 \end{aligned} \tag{28}$$

The exact forms from this equation are eliminated with the help of Stoke's integral theorem. As a consequence of Hamilton Principle, one gets two sets equations of gravitational field

$$\frac{1}{2}\Omega^{kl} \wedge \eta_{kli} - \Lambda\eta_i + 8\pi t_i = 0 \iff \frac{1}{2}\Omega^{kl} \wedge \eta_{kli} - \Lambda\eta_i = -8\pi t_i, \tag{29}$$

$$\frac{1}{8\pi}D\eta^j_i + s_i{}^j = 0 \iff D\eta^j_i = -8\pi s_i{}^j \tag{30}$$

and

$$L_A = 0. \tag{31}$$

$L_A = 0$ represent equations of motion for matter field. These equations are not intrinsic in further our considerations, so we will omit them. We are interested only in the gravitational field equations which are given by the equations (29)-(30).

In vacuum where $t_i = s_i{}^j = 0 \implies D\eta_i{}^j = 0$ and we get the standard vacuum Einstein's equations (EE) with cosmological constant Λ and pseudoriemannian geometry

$$\frac{1}{2}\Omega^{kl} \wedge \eta_{kli} - \Lambda\eta_i = 0. \tag{32}$$

In general, we have the Einstein-Cartan equations and Riemann-Cartan geometry (a metric geometry with torsion).
The standard GR we obtain also if we put $D\eta_i{}^k = 0 \implies s_i{}^k = 0$, i.e., if we confine to pseudoriemannian geometry. It is GR inside spinless matter with equations

$$\frac{1}{2}\Omega^{kl} \wedge \eta_{kli} - \Lambda\eta_i = -8\pi t_i. \tag{33}$$

One can show that $\frac{1}{2}\Omega^{kl} \wedge \eta_{kli} = -G_i^s\eta_s$ and $t_i = T_i{}^s\eta_s$, where the Einstein tensor G_i^s is defined as follows

$$G_i^s = R_i^s - \frac{1}{2}\delta_i^s R. \tag{34}$$

Moreover, if $s_i{}^k = 0$ then the matter tensor $T_i{}^s$, defined by the decomposition $t_i = T_i{}^s\eta_s$, is symmetric, i.e. $T^{ik} = T^{ki}$. Because $\Lambda\eta_i = \Lambda\delta_i^s\eta_s$, then we get from (33)

$$-G_i^s\eta_s - \Lambda\delta_i^s\eta_s = -8\pi T_i{}^s\eta_s. \tag{35}$$

Multiplying this equation by -1 and omitting η_s, we can see that there are Einstein equations with cosmological constant Λ in tensorial notation

$$G_i^s + \Lambda\delta_i^s = 8\pi T_i{}^s. \tag{36}$$

2 Einstein-Palatini action integral for General Relativity in vacuum and with positive cosmological constant Λ as integral action for a gauge field

Now, getting back to Einstein-Palatini action in vacuum

$$\begin{aligned} S_{E-P} &= \frac{1}{4\kappa}\int_{\mathcal{D}}\Big(\vartheta^i \wedge \vartheta^j \wedge \Omega^{kl} - \frac{\Lambda}{6}\vartheta^i \wedge \vartheta^j \wedge \vartheta^k \wedge \vartheta^l\Big)\eta_{ijkl} \\ &= \frac{1}{4\kappa}\int_{\mathcal{D}}\Big(\vartheta^i \wedge \vartheta^j \wedge \Omega^{kl}\eta_{ijkl} - \frac{\Lambda}{6}\vartheta^i \wedge \vartheta^j \wedge \vartheta^k \wedge \vartheta^l\eta_{ijkl}\Big) \end{aligned} \tag{37}$$

and using the definition of the duality operator $\star$ [1]

$$\star := -\frac{\eta_{ijkl}}{2} \qquad \implies \eta_{ijkl} = -2\star \tag{38}$$

one gets

$$\eta_{ijkl}\Omega^{kl} = -2 \star \Omega_{ij}, \tag{39}$$

$$\eta_{ijkl}\vartheta^k \wedge \vartheta^l = -2 \star \big(\vartheta_i \wedge \vartheta_j\big). \tag{40}$$

Thus the Einstein-Palatini action has the following form

$$S_{E-P} = -\frac{1}{2\kappa}\int\limits_{\mathcal{D}}\Big(\vartheta^i\wedge\vartheta^j\wedge\star\Omega_{ij} - \frac{\Lambda}{6}\vartheta^i\wedge\vartheta^j\wedge\star\big(\vartheta_i\wedge\vartheta_j\big)\Big)$$
$$= -\frac{1}{2\kappa}\int\limits_{\mathcal{D}} tr\Big(\vartheta\wedge\vartheta\wedge\star\Omega - \frac{\Lambda}{6}\vartheta\wedge\vartheta\wedge\star\big(\vartheta\wedge\vartheta\big)\Big). \tag{41}$$

Let us introduce the corrected curvature $\hat{F}$

$$\hat{F} := \Omega - \frac{\Lambda}{3}\vartheta\wedge\vartheta \quad \Longrightarrow\ \vartheta\wedge\vartheta = \frac{3}{\Lambda}\Big(\Omega - \hat{F}\Big). \tag{42}$$

Substituting the last formula into Einstein-Palatini action we get

$$S_{E-P} = -\frac{1}{2\kappa}\int\limits_{\mathcal{D}} tr\Big(\vartheta\wedge\vartheta\wedge\star\Omega - \frac{\Lambda}{6}\vartheta\wedge\vartheta\wedge\star\big(\vartheta\wedge\vartheta\big)\Big)$$
$$= -\frac{1}{2\kappa}\int\limits_{\mathcal{D}} tr\Big[\frac{3}{\Lambda}\big(\Omega-\hat{F}\big)\wedge\star\Omega - \frac{\Lambda}{6}\frac{9}{\Lambda^2}\big(\Omega-\hat{F}\big)\wedge\star\big(\Omega-\hat{F}\big)\Big]$$
$$= -\frac{3}{4\Lambda\kappa}\int\limits_{\mathcal{D}} tr\Big(2\big(\Omega-\hat{F}\big)\wedge\star\Omega - \big(\Omega-\hat{F}\big)\wedge\star\big(\Omega-\hat{F}\big)\Big)$$
$$= -\frac{3}{4\Lambda\kappa}\int\limits_{\mathcal{D}} tr\Big[2\Omega\wedge\star\Omega - 2\hat{F}\wedge\star\Omega - \Omega\wedge\star\Omega + \hat{F}\wedge\star\Omega + \Omega\wedge\star\hat{F} - \hat{F}\wedge\star\hat{F}\Big]$$
$$= -\frac{3}{4\Lambda\kappa}\int\limits_{\mathcal{D}} tr\Big[\Omega\wedge\star\Omega - \hat{F}\wedge\star\Omega + \Omega\wedge\star\hat{F} - \hat{F}\wedge\star\hat{F}\Big] \tag{43}$$

Because $-\hat{F}\wedge\star\Omega + \Omega\wedge\star\hat{F}$ reduces, then we finally have

$$S_{E-P} = -\frac{3}{4\Lambda\kappa}\int\limits_{\mathcal{D}} tr\Big[\Omega\wedge\star\Omega - \hat{F}\wedge\star\hat{F}\Big]. \tag{44}$$

The expression $tr\big(\Omega\wedge\star\Omega\big) = \eta_{ijkl}\Omega^{ij}\wedge\Omega^{kl}$ is in four dimensions a topological invariant called Euler's form, which does not influence the equations of motion [12]. Hence, in 4-dimensions the Einstein-Palatini action is equivalent to

$$S_{E-P} = \frac{3}{4\Lambda\kappa}\int\limits_{\mathcal{D}} tr\Big(\hat{F}\wedge\star\hat{F}\Big). \tag{45}$$

We see that the Einstein-Palatini action in 4-dimensions is efectively the functional which is quadratic function of the corrected Riemannian curvature, i.e., it has a form of the action for a gauge field.

Only one difference is that in (45) we have the star operator $\star$, which is different from Hodge star operator. Namely, our star operator acts onto "interior" indices (tetrad's indices), not onto forms as Hode duality operator does [2, 12].

3 Geometrical interpretation of the corrected curvature $\hat{F}$

Let $P(M_4, GdS)$ denotes the principal bundle of de Sitter basis over a manifold M_4 (spacetime) with de Sitter group(GdS) [5, 13] as a structure group. This group is isomorphic to the group $SO(4,1)$ [3, 5, 13]. Let $\widetilde{\omega}$ be 1-form of connection in the principle fibre bundle $P(M_4, GdS)$. The form $\widetilde{\omega}$ has values in the algebra $AlgGdS$ of the group GdS. This algebra splits (as a vector space) into direct sum

$$AlgGdS = so(3,1) \oplus R^{(3,1)}. \tag{46}$$

$so(3,1)$ denotes here algebra of the group SO(3, 1), which is isomorphic to Lorentz group $\mathcal{L}$, and $R^{(3,1)}$ is a 4-dimensional vector space of generalised translations (translations in the curved spacetime). This vector space we identify with the de Sitter spacetime [1, 2, 13].

Let $P(M_4, \mathcal{L})$ denotes the principal bundle of Lorentz basis over the manifold M_4. There exists a morphism of principal bundles [4]

$$f : P(M_4, \mathcal{L}) \longrightarrow P(M_4, GdS). \tag{47}$$

This morphism creates pull-back $f_*\widetilde{\omega}$ of the form $\widetilde{\omega}$ onto the bundle $P(M_4, \mathcal{L})$. Let us denote this pull-back by A. A is a 1-form on $P(M_4, \mathcal{L})$ with values in the direct sum [4]

$$so(3,1) \oplus R^{(3,1)}. \tag{48}$$

Hence, we have a natural decomposition [4]

$$A = f_*\widetilde{\omega} = \omega + \theta, \tag{49}$$

where ω is a 1-form on $P(M_4, \mathcal{L})$ with values in the algebra $so(3,1)$ and θ is a 1-form on $P(M_4, \mathcal{L})$ with values on $R^{(3,1)}$. ω is a connection on the bundle $P(M_4, \mathcal{L})$.

Let us compute a 2-form curvature $\hat{F}$ of the pulled back A. From the definition we have

$$\begin{aligned}
\hat{F} &= dA + \frac{1}{2}\Big[A, A\Big] \\
&= d\big(\omega + \theta\big) + \frac{1}{2}\Big[\omega + \theta, \omega + \theta\Big] \\
&= d\omega + \frac{1}{2}\Big[\omega, \omega\Big] + d\theta + \frac{1}{2}\Big[\omega, \theta\Big] + \frac{1}{2}\Big[\theta, \omega\Big] + \frac{1}{2}\Big[\theta, \theta\Big].
\end{aligned} \tag{50}$$

We are going to introduce to our equations bases $\widetilde{M}_{ik} = -\widetilde{M}_{ki}$ of algebra $so(3,1)$ and e_i of vector space $R^{(3,1)}$. In these bases, we have

$$\omega = \omega^i{}_k \widetilde{M}^k_i = \omega^{ik}\widetilde{M}_{ki},\ \theta = \theta^i e_i. \tag{51}$$

$(\widetilde{M}_{ik}, e_l)$ form together the algebra of the de Sitter group (the basis of algebra AlgdS). Our elements $\widetilde{M}_{ik} = -\widetilde{M}_{ki}$ are real and connected with elements $M_{ik} = M_{ki}$ used in [13] in the following way

$$\widetilde{M}_{ik} = \frac{i}{2} M_{ik} \Rightarrow M_{ik} = -2i\widetilde{M}_{ik}. \tag{52}$$

In the terms of the elements $[\widetilde{M}_{ik}, e_l]$ the comutational relations for the algebra $so(4,1) = so(3,1) \oplus R^{(3,1)}$ read

$$[\widetilde{M}_{ij}, \widetilde{M}_{kl}] = \frac{1}{2}\big(\eta_{ik}\widetilde{M}_{jl} + \eta_{jl}\widetilde{M}_{ik} - \eta_{il}\widetilde{M}_{jk} - \eta_{jk}\widetilde{M}_{il}\big) \tag{53}$$

$$[e_i, \widetilde{M}_{jk}] = \frac{1}{2}\big(\eta_{ik}e_j - \eta_{ij}e_k\big) \tag{54}$$

$$[e_i, e_j] = \frac{2\widetilde{M}_{ij}}{R^2} \tag{55}$$

The following commutation relations are important in the further considerations [5, 6, 13].

$$[\widetilde{M}_{ki}, e_l] = \frac{1}{2}\big(\eta_{kl}e_i - \eta_{il}e_k\big) \tag{56}$$

$$[e_i, e_k] = \frac{2\widetilde{M}_{ik}}{R^2}, \tag{57}$$

where R is the radius of the de Sitter spacetime. Using the above equations we have

$$\hat{F} = \Omega_\omega + \frac{1}{2}\omega^i{}_k \wedge \theta^l \Big[\widetilde{M}^k{}_i, e_l\Big] + \frac{1}{2}\theta^l \wedge \omega^i{}_k \Big[e_l, \widetilde{M}^k{}_i\Big] + \frac{1}{2}\theta^i \wedge \theta^k \Big[e_i, e_k\Big] + d\theta^i e_i, \tag{58}$$

where $\Omega_\omega = d\omega + \frac{1}{2}[\omega, \omega]$ is the curvature 2-form of the connection's ω. Taking into consideration the commutation relations in algebra AlgdS given by the formulas (56) and (57), we obtain

$$\begin{aligned}
\hat{F} &= \Omega_\omega + \omega^i{}_k \wedge \theta^l \Big[\widetilde{M}^k{}_i, e_l\Big] + \theta^i \wedge \theta^k \frac{\widetilde{M}_{ik}}{R^2} + d\theta^i e_i \\
&= \Omega_\omega^{ik}\widetilde{M}_{ki} + \big(\omega^i{}_{.k} \wedge \theta^k\big)e_i + d\big(\theta^i\big)e_i - \frac{\theta^i \wedge \theta^k \widetilde{M}_{ki}}{R^2} \\
&= \Omega_\omega^{ik}\widetilde{M}_{ki} - \frac{\theta^i \wedge \theta^k \widetilde{M}_{ki}}{R^2} + \big(d\theta^i + \omega^i{}_{.k} \wedge \theta^k\big)e_i \\
&= \Omega_{corr}^{ik}\widetilde{M}_{ki} + \big(\mathcal{D}_\omega\theta^i\big)e_i \\
&= \Omega_{corr}^{ik}\widetilde{M}_{ki} + \Theta^i e_i
\end{aligned} \tag{59}$$

$\Omega_{corr} := \Omega_\omega - \frac{\theta\wedge\theta}{R^2} = \Omega_\omega - \frac{\Lambda}{3}\theta \wedge \theta$ and it denotes the corrected curvature of the connection ω on the bundle $P(M_4, \mathcal{L})$ and $\Theta = \mathcal{D}_\omega\theta$ is a torsion of the connection ω.
If we adjust the connection $\widetilde{\omega}$ in such a way that the connection ω is torsionless ($\Theta = 0$), i.e. if ω is Levi-Civita connection, then we get (after leaving the basis $so(3,1)$ and $R^{(3,1)}$)

$$\hat{F} = \Omega_\omega - \frac{\Lambda}{3}\theta \wedge \theta = \Omega_{corr}\ . \tag{60}$$

In the Section 2 we gave the definition of the corrected curvature F as follows:

$$\hat{F} := \Omega - \frac{\Lambda}{3}\vartheta \wedge \vartheta. \tag{61}$$

As one can see this curvature is a curvature of the connection

$$A := f_*\widetilde{\omega} = \omega + \Theta \tag{62}$$

if $\Theta = 0$.

4 Conclusion

In this article we have shown that in four dimentions the action integral for GR with a positive cosmological constant Λ can be written in an analogical form to the form of the action integral for the typical gauge field. However, there is one difference - the star. Instead of the Hodge star, we have slightly different star called the duality operator [2, 12].
Our result is important because it shows that there is no need to generalize GR and construct very complicated gravitational theories to obtain a gravitational theory as a gauge theory. The ordinary GR formulated in terms of tetrads and spin connection with cosmological constant $\Lambda > 0$ is already a gauge theory.
Some scientists [1, 2, 3] were concerned with this problem and they came to the similar conclusions as ours, but they applied in their works the Cartan's approach to the connection in the principal bundle [2, 13, 14]. This approach is not well known among geometrists and relativists. We have used only the standard theory of connection in the principal bundle which was created by Ehresmann - Cartan's student [4, 8]. His approach is commonly used in differential geometry and in relativity. We would like to emphasize that the formulation of the EP action in the form (45) can be important for quantizing of general relativity (because gauge fields are quantized).

Acknowledgements

One of the authors (M.D.) would like to thank J. Garecki for consultative help and constructive suggestion on matter in this article.

Appendix 1

η forms and operations with them [11]

Following [11] we define

$$\eta_{ijkl} = \sqrt{|g|}\epsilon_{ijkl} \tag{A.1}$$

where ϵ_{ijkl} is Levi-Civita pseudotensor with properties

$$\epsilon_{ijkl} = \begin{cases} 1 & \text{if the sequence of indices ijkl is an even permutation} \\ & \text{of the sequence 0, 1, 2, 3;} \\ -1 & \text{if it is an odd permutation;} \\ 0 & \text{if the sequence of indices ijkl is not an even permutation} \\ & \text{of the sequence 0, 1, 2, 3} \end{cases} . \tag{A.2}$$

and we take $\eta_{0123} = \sqrt{|g|}$. In Lorentzian coreper $|g| = 1$.

One has [11]

$$\eta_{ijk} = \vartheta^l \wedge \eta_{ijkl} \tag{A.3}$$

$$\eta_{ij} = \frac{1}{2}\vartheta^k \wedge \eta_{ijk} \tag{A.4}$$

$$\eta_i = \frac{1}{3}\vartheta^j \wedge \eta_{ij} \tag{A.5}$$

$$\eta = \frac{1}{4}\vartheta^i \wedge \eta_i \tag{A.6}$$

$$\vartheta^n \wedge \eta_{kli} = \delta^n_i \eta_{kl} + \delta^n_l \eta_{ik} + \delta^n_k \eta_{li} \tag{A.7}$$

$$\vartheta^m \wedge \eta_{kl} = \delta^m_l \eta_k - \delta^m_k \eta_l \tag{A.8}$$

$$\vartheta^j \wedge \eta_i = \delta^j_i \eta \tag{A.9}$$

The forms η, η_i, η_{ij}, η_{ijk} are Hodge dual to the forms 1, ϑ^i, $\vartheta^i \wedge \vartheta^j$, $\vartheta^i \wedge \vartheta^j \wedge \vartheta^k$ respectively [11].

References

[1] D. K. Wise, *"MacDowell-Mansouri Gravuty and Cartan Geometry"*, CQG, **27** (2010) 155010 (arXiv:gr-qc/0611154v2, 15 May 2009)

[2] D. K. Wise, *"Symmetric Space Cartan Connections and Gravity in Three and Four Dimensions"*, arXiv:0904.1738v2 [math.DG], 3 August 2009

[3] A. Randomo, *"Gauge Gravity: a forward-looking introduction"*, arXiv: 1010.5822v1 [gr-qc], 27 October 2010

[4] S. Kobayashi, K. Nomizu, *"Foundations of Differential Geometry"*, Interscience Publishers, a division of John Wiley and Sons, New York , London 1963

[5] F. Gürsey, *"Introduction to Group Theory" an article in "Groups and Topology in Relativity"*,C. DeWitt and B. DeWitt (editors), Gordon and Breach, London 1964

[6] A. Dubničkova, *"Topological Groups for Physicists"*, Dubna 1987 (in Russian)

[7] J. Mozrzymas, *"Applications of Group Theory in Modern Physics"*, National Scientific Publishers PWN, Wrocław 1967 (in Polish)

[8] J. Gancarzewicz, *"Foundations of Modern Differential Geometry"*, SCRIPT, Warsaw 2010 (in Polish)

[9] R. Sulanke, P. Wintgen, *"Differentialgeometrie und Faserbuendel"*, Copyright by VEB Deutscher Verlag der Wissenschaften, Berlin 1972

[10] W. Kopczyński, A. Trautman, *"Spacetime and Gravitation"*, National Scientific Publishers PWN, Warszawa 1984 (in Polish - there exists English translation)

[11] A. Trautman, *"Einstein-Cartan Theory"*, Symposia Mathematica, **12** (1973) 139

[12] K. Hayashi, T. Shirafuji, *"Gravity from Poincare Gauge Theory of the Fundamental Particles. Part V" an article in "Progress of Theoretical Physics"*, **65** (1981) 525

[13] W. Drechsler, M.E. Mayer, *"Fiber Bundle Techniques in Gauge Theories"* , an article in ""Lectures Notes in Physics" Vol.67, Springer-Verlag, Berlin · Heidelberg · New York 1977

[14] R. W. Sharpe, *"Differential Geometry. Cartan's Generalization of Klein's Erlangen Program"* , Springer-Verlag, New York · Berlin · Heidelberg 2000